



SCN4A gene

sodium voltage-gated channel alpha subunit 4

Normal Function

The *SCN4A* gene belongs to a family of genes that provide instructions for making sodium channels. These channels, which transport positively charged sodium atoms (sodium ions) into cells, play a key role in a cell's ability to generate and transmit electrical signals.

The *SCN4A* gene provides instructions for making a critical part (the alpha subunit) of sodium channels that are abundant in muscles used for movement (skeletal muscles). For the body to move normally, these muscles must tense (contract) and relax in a coordinated way. One of the changes that helps trigger muscle contractions is the flow of positively charged atoms (ions), including sodium, into muscle cells. Channels made with the *SCN4A* protein control the flow of sodium ions into these cells.

Health Conditions Related to Genetic Changes

congenital myasthenic syndrome

hyperkalemic periodic paralysis

More than 10 mutations in the *SCN4A* gene have been found to cause hyperkalemic periodic paralysis, a condition that causes episodes of extreme muscle weakness that are often associated with high levels of potassium in the blood (hyperkalemia). The mutations change single building blocks (amino acids) in the *SCN4A* protein, which alters the structure and function of sodium channels in skeletal muscle cells. These changes delay the closing of channels made with the *SCN4A* protein or prevent the channels from staying closed. As a result, sodium ions continue flowing into muscle cells abnormally. This increase in sodium ions triggers the release of potassium from muscle cells, which causes more sodium channels to open and stimulates the flow of even more sodium ions into these cells. These changes in ion transport reduce the ability of skeletal muscles to contract, leading to episodes of muscle weakness or paralysis.

hypokalemic periodic paralysis

At least seven mutations in the *SCN4A* gene have been identified in people with hypokalemic periodic paralysis, a condition that causes episodes of extreme muscle weakness that are associated with low levels of potassium in the blood (hypokalemia). Mutations in the *SCN4A* gene account for about 10 percent of all

cases of this condition. Each of the known mutations changes a single amino acid in the *SCN4A* protein, which alters the structure and function of sodium channels in skeletal muscle cells. The abnormal channels change the normal flow of sodium ions, which prevents muscles from contracting normally. Low potassium levels also contribute to this problem. Because muscle contraction is needed for movement, these changes in ion transport lead to long-lasting episodes of severe muscle weakness.

paramyotonia congenita

At least 18 mutations in the *SCN4A* gene are known to cause paramyotonia congenita, a muscle disease characterized by episodes of sustained muscle tensing (myotonia) that prevent muscles from relaxing normally. The *SCN4A* gene mutations that cause this condition each change a single amino acid in the *SCN4A* protein, which alters the structure and function of sodium channels in skeletal muscle cells. The most common genetic changes replace the amino acid arginine with one of several other amino acids at protein position 1448.

Mutations delay the closing of channels made with the *SCN4A* protein and, once the channels are closed, cause them to open again too quickly. These changes increase the flow of sodium ions into skeletal muscle cells. An influx of extra sodium ions triggers prolonged muscle contractions, which underlie the episodes of myotonia characteristic of paramyotonia congenita. Muscles with sustained high levels of sodium ions may become unable to contract at all, resulting in attacks of muscle weakness.

The effects of *SCN4A* gene mutations on the altered ion channels may be increased by cold temperatures, which may help explain why signs and symptoms can be triggered by exposure to cold.

potassium-aggravated myotonia

Several mutations in the *SCN4A* gene result in potassium-aggravated myotonia, a condition that causes episodes of myotonia that prevent muscles from relaxing normally. The resulting muscle stiffness may be aggravated by eating potassium-rich foods. The most common genetic changes associated with potassium-aggravated myotonia replace the amino acid glycine with one of several other amino acids at protein position 1306. These mutations delay the closing of channels made with the *SCN4A* protein, which increases the flow of sodium ions into skeletal muscle cells. The presence of excess potassium stimulates the flow of even more sodium ions into these cells. These changes in ion transport trigger prolonged muscle contractions, which underlie the muscle stiffness characteristic of potassium-aggravated myotonia.

other disorders


A mutation in the *SCN4A* gene is also responsible for one form of congenital myasthenic syndrome, a muscle disorder that appears shortly after birth. People

with this condition have general muscle weakness and recurrent attacks of paralysis that specifically affect muscles used for speaking and breathing. The *SCN4A* gene mutation associated with this condition replaces the amino acid valine with the amino acid glutamic acid at protein position 1442 (written as Val1442Glu or V1442E). This genetic change alters the structure and function of sodium channels in skeletal muscle cells. The channels stay closed abnormally after repeated muscle contractions, reducing the flow of sodium ions into muscle cells. This disruption in ion transport reduces further muscle contraction, leading to muscle weakness and episodes of paralysis.

Chromosomal Location

Cytogenetic Location: 17q23.3, which is the long (q) arm of chromosome 17 at position 23.3

Molecular Location: base pairs 63,938,554 to 63,972,918 on chromosome 17 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- Na(V)1.4
- Nav1.4
- SCN4A_HUMAN
- skeletal muscle voltage-dependent sodium channel type IV alpha subunit
- SkM1
- sodium channel, voltage gated, type IV alpha subunit
- sodium channel, voltage-gated, type IV, alpha
- sodium channel, voltage-gated, type IV, alpha subunit
- voltage-gated sodium channel type 4 alpha

Additional Information & Resources

Educational Resources

- Basic Neurochemistry (sixth edition, 1999): Na⁺ Channel Mutations Cause Periodic Paralysis
<https://www.ncbi.nlm.nih.gov/books/NBK28162/#A3038>
- Biochemistry (fifth edition, 2002): Specific Channels Can Rapidly Transport Ions Across Membranes
<https://www.ncbi.nlm.nih.gov/books/NBK22509/>

GeneReviews

- Congenital Myasthenic Syndromes
<https://www.ncbi.nlm.nih.gov/books/NBK1168>
- Hyperkalemic Periodic Paralysis
<https://www.ncbi.nlm.nih.gov/books/NBK1496>
- Hypokalemic Periodic Paralysis
<https://www.ncbi.nlm.nih.gov/books/NBK1338>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28SCN4A%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- SODIUM CHANNEL, VOLTAGE-GATED, TYPE IV, ALPHA SUBUNIT
<http://omim.org/entry/603967>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_SCN4A.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=SCN4A%5Bgene%5D>
- HGNC Gene Family: Sodium voltage-gated channel alpha subunits
<http://www.genenames.org/cgi-bin/genefamilies/set/1203>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=10591

- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/6329>
- UniProt
<http://www.uniprot.org/uniprot/P35499>

Sources for This Summary

- Burge JA, Hanna MG. Novel insights into the pathomechanisms of skeletal muscle channelopathies. *Curr Neurol Neurosci Rep*. 2012 Feb;12(1):62-9. doi: 10.1007/s11910-011-0238-3. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22083238>
- Carle T, Lhuillier L, Luce S, Sternberg D, Devuyst O, Fontaine B, Tabti N. Gating defects of a novel Na⁺ channel mutant causing hypokalemic periodic paralysis. *Biochem Biophys Res Commun*. 2006 Sep 22;348(2):653-61. Epub 2006 Jul 28.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16890191>
- Colding-Jørgensen E, Duno M, Vissing J. Autosomal dominant monosymptomatic myotonia permanens. *Neurology*. 2006 Jul 11;67(1):153-5.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16832098>
- Dice MS, Abbruzzese JL, Wheeler JT, Groome JR, Fujimoto E, Ruben PC. Temperature-sensitive defects in paramyotonia congenita mutants R1448C and T1313M. *Muscle Nerve*. 2004 Sep;30(3):277-88.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15318338>
- GeneReview: Hyperkalemic Periodic Paralysis
<https://www.ncbi.nlm.nih.gov/books/NBK1496>
- GeneReview: Hypokalemic Periodic Paralysis
<https://www.ncbi.nlm.nih.gov/books/NBK1338>
- Groome JR, Fujimoto E, Ruben PC. K-aggravated myotonia mutations at residue G1306 differentially alter deactivation gating of human skeletal muscle sodium channels. *Cell Mol Neurobiol*. 2005 Nov;25(7):1075-92.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16392038>
- Hantaï D, Richard P, Koenig J, Eymard B. Congenital myasthenic syndromes. *Curr Opin Neurol*. 2004 Oct;17(5):539-51. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15367858>
- Jurkat-Rott K, Holzherr B, Fauler M, Lehmann-Horn F. Sodium channelopathies of skeletal muscle result from gain or loss of function. *Pflugers Arch*. 2010 Jul;460(2):239-48. doi: 10.1007/s00424-010-0814-4. Epub 2010 Mar 17. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20237798>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2883924/>
- Jurkat-Rott K, Weber MA, Fauler M, Guo XH, Holzherr BD, Paczulla A, Nordsborg N, Joechle W, Lehmann-Horn F. K⁺-dependent paradoxical membrane depolarization and Na⁺ overload, major and reversible contributors to weakness by ion channel leaks. *Proc Natl Acad Sci U S A*. 2009 Mar 10;106(10):4036-41. doi: 10.1073/pnas.0811277106. Epub 2009 Feb 18.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19225109>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2644652/>

- Miller TM, Dias da Silva MR, Miller HA, Kwiecinski H, Mendell JR, Tawil R, McManis P, Griggs RC, Angelini C, Servidei S, Petajan J, Dalakas MC, Ranum LP, Fu YH, Ptáček LJ. Correlating phenotype and genotype in the periodic paralyses. *Neurology*. 2004 Nov 9;63(9):1647-55.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15534250>
- Tamaoka A. Paramyotonia congenita and skeletal sodium channelopathy. *Intern Med*. 2003 Sep; 42(9):769-70.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14518660>
- Tsujino A, Maertens C, Ohno K, Shen XM, Fukuda T, Harper CM, Cannon SC, Engel AG. Myasthenic syndrome caused by mutation of the SCN4A sodium channel. *Proc Natl Acad Sci U S A*. 2003 Jun 10;100(12):7377-82. Epub 2003 May 23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12766226>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC165883/>
- Vicart S, Sternberg D, Fontaine B, Meola G. Human skeletal muscle sodium channelopathies. *Neurol Sci*. 2005 Oct;26(4):194-202. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16193245>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/SCN4A>

Reviewed: August 2013
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services